

# STAT<sup>+</sup>CUP



## STATCUP<sup>®</sup>

The STATCUP<sup>®</sup> is a fully integrated on-site urine drug test cup. Simply collect the sample and read the results. It's that easy! See back for full procedure details and how to read results.

- Fast
- Accurate
- Reliable
- Easy-to-Use
- FDA approved
- Affordable

**CALL FOR A  
FREE SAMPLE!**

### TEST FEATURES

- No Urine Exposure
- No Steps to Activate
- Screw Top Lid
- Temperature Strip
- Test Up to 15 Drugs (see below)
- Available with Built-in Adulterants
- Transport Sample to Lab

### Drugs of Abuse:

Marijuana  
Cocaine  
Opiates  
Methamphetamine  
Amphetamine  
Phencyclidine  
Benzodiazepines  
Barbiturates  
Methadone  
Ecstasy  
Tricyclic Antidepressants  
Oxycodone  
Propoxyphene\*  
Buprenorphine\*  
6-Monoacetylmorphine\*  
Alcohol\*

### Adulterants:

pH (acidic additives)  
Specific Gravity (sample dilution)  
Oxidants (bleach/oxidizing agents)

\*For Forensic Use Only



**Toll Free: 866-933-0964**  
[www.micro-distributing.com](http://www.micro-distributing.com)



CALL FOR A FREE SAMPLE!

# STAT CUP

## EASY-TO-USE FAST ACCURATE

### PROCEDURE

**1** Remove Cap.





**2** Donor provides specimen.




**3** Replace the Cap securely. Peel off the blue label that covers the result window.



**4** Read results at 5 minutes. Test results remain stable for a minimum of 60 minutes.



### INTERPRETING RESULTS




Interpret adulterant test strip between 3-5 minutes. Refer to enclosed color chart for an accurate interpretation.

Toll Free: 866-933-0964

Phone: 254-939-8923

Fax: 254-939-5867

sales@micro-distributing.com



All positive results are presumptive and should be confirmed by an alternative method (e.g. GC/MS). A photocopy of the results can be made by placing the entire sealed cup with the card face down on the copier.